

# Efficacy of Diclofenac Rectal Suppository in Patients with Laparoscopic Cholecystectomy: A Prospective Randomized Double Blinded Clinical Trial

Mohammadreza Arab<sup>1</sup>, Hossein Saeedi Motahhar<sup>2</sup>, Abdolreza Pazouki<sup>2,\*</sup>, Zeinab Tamannaie<sup>2</sup>, Babak Arabpour<sup>2</sup>, Hossein Fahimi<sup>2</sup>

<sup>1</sup>Department of Surgery, Firoozgar Hospital, Minimally Invasive Surgery Research Center, Iran University of Medical Sciences, Tehran, IR Iran

<sup>2</sup>Minimally Invasive Surgery Research Center, Iran University of Medical Sciences, Tehran, IR Iran

\*Corresponding author: Abdolreza Pazouki, Minimally Invasive Surgery Research Center, Iran University of Medical Sciences, Tehran, IR Iran., Tel.: +98-2166555447, Fax: +98-2166501113, E-mail: apazouki@yahoo.com.

Received: May 13, 2013; Accepted: July 14, 2013

**Background:** Post laparoscopic cholecystectomy pain management can reduce recovery and discharge time. Non-steroidal anti-inflammatory drugs and opioids are used for this purpose.

**Objectives:** This randomized clinical trial evaluates the efficacy of diclofenac rectal suppository for the management of postoperative pain.

**Patients and Methods:** Forty four patients were randomized to receive either 100 mg diclofenac rectal suppository or placebo at the time of recovery and three hours later after laparoscopic cholecystectomy. Postoperative visual analogue pain scale (VAS, ranges 0 to 10 cm) and adverse reactions were recorded over a 24-hour period. If VAS score was  $\geq 7$ , 25mg, pethedin was given intravenously as a rescue analgesic.

**Results:** In both groups, VAS score was reduced in 24 hours. It was statistically lower in diclofenac group rather than placebo group in all intervals except at the time of recovery. Moreover, the mean pethedin consumption dose and the incidence of administration of postoperative rescue analgesic were statistically lower in diclofenac group. Postoperative bleeding was not statistically different between two groups.

**Conclusions:** Diclofenac rectal suppository provided simple and safe pain relief in laparoscopic cholecystectomy.

**Keywords:** Cholecystectomy, Laparoscopic; Diclofenac; Analgesics, Opioid

## 1. Background

Although minimally invasive surgeries have reduced the postoperative pain, most of the patients still require opioids after surgery. Opioids have important side effects including respiratory depression, drowsiness, apnea, nausea, vomiting, and ileus (1, 2). Non-steroidal Anti-inflammatory Drugs (NSAIDs) are safer than opioids with the similar effect on postoperative pain (2). It has been shown that using intramuscular or intravenous diclofenac and ketorolac in upper abdominal and gynecologic surgeries could decrease the need for opioids. Muscle and kidney damage and increasing the risk of bleeding are important side effects of intramuscular administration of diclofenac (3).

## 2. Objectives

This study was done to evaluate the effect of rectal suppository diclofenac on post laparoscopic cholecystectomy pain.

## 3. Patients and Methods

The protocol of this prospective randomized double-blinded, controlled clinical trial was approved by the local ethical committee of the medical faculty, Tehran University of Medical Sciences. Forty four patients candidate for laparoscopic cholecystectomy of gallbladder stone enrolled in this study from January 2011 to January 2013 at Rasool Akram and Firouzgar hospital, Tehran, Iran., . Exclusion criteria were acute cholecystitis, patient refusal to enter or continue the study, morbid obesity (body mass index  $\geq 30$ ), chemical substance abuse, chronic or recent use of analgesics, chronic pain, a history of significant cardiopulmonary, hepatic or renal disease, hypersensitivity to the medications, any contraindication of non-opioid analgesics, asthma, peptic ulcer, upper or lower gastrointestinal bleeding, pregnancy, and breastfeeding. The patients who required analgesic the night before surgery were also excluded from this study. A written informed consent was obtained from all patients and

### Implication for health policy/practice/research/medical education:

Post laparoscopic cholecystectomy pain management can reduce recovery and discharge time. Non-steroidal anti-inflammatory drugs and opioids are used for this purpose. This randomized clinical trial evaluates the efficacy of diclofenac rectal suppository in the management of postoperative pain.

Copyright © 2013, Minimally Invasive Surgery Research Center and Mediterranean. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

they were randomly assigned into either treatment (100 mg diclofenac rectal suppository at recovery time and 3 hours later) or placebo group (using placebo at recovery time and 3 hours later) using computer table of random numbers. All patients underwent pain assessment with Visual Analogue Scale (VAS, ranges 0 – 10 cm) at the time of recovery, and 3, 6, 12, and 24 hours after operation. Patients were given 25 mg Pethedin intravenously if their VAS score was  $\geq 7$ . Pain scores, total Pethedin consumption, patients requests for administration of postoperative analgesics, and also intraoperative and postoperative complications including bleeding, visceral injury, nausea, vomiting, and headache were recorded.

### 3.1. Statistical Analysis

Continuous variables are summarized by mean  $\pm$  standard deviation and categorical variables are expressed as percentage (%). Univariate analyses were performed by Chi square, student T-test, Mann-Whitney U test using SPSS software (version 17; SPSS Inc, Chicago, IL). Results were considered significant if P values were less than 0.05.

## 4. Results

Baseline characteristics of patients were not statistically different between two groups (Table 1). One patient (4.5%) in diclofenac group and another one (4.5%) in placebo group underwent Magnetic Resonant Cholangiopancreatography (MRCP). Moreover, six patients (27%) in diclofenac group and eight patients (36%) in placebo group underwent Endoscopic Retrograde Cholangiopancreatography (ERCP) before laparoscopic cholecystectomy. Intraoperative findings including gallbladder adhesion was detected in 17 patients (77.3%) of diclofenac group and 12 patients (54.5%) of placebo group which was not statistically different among them ( $P = 0.101$ ). Moreover, three patients in diclofenac group (13.6%) and six patients (27.3%) in placebo group had intraoperative bleeding which was not statistically different, too ( $P = 0.228$ ). Five patients (22.7%) of diclofenac group and 12 patients (54.5%) of placebo group required analgesic treatment for one day. Furthermore, one patient (4.5%) of diclofenac group and nine patients (40.9%) of placebo group required analgesic treatment for two days. Results showed that the incidence for administration of postoperative analgesic in diclofenac group was statistically lower than that in placebo group ( $P < 0.001$ ). Similarly, the mean Pethedin consumption dose in diclofenac group ( $18.18 \pm 15.77$  mg) was statistically lower than that in placebo group ( $66.59 \pm 18.80$  mg) with  $P$  value  $< 0.001$ . Table 2 demonstrates the mean VAS score at the time of recovery, 3, 6, 12, and 24 hours after cholecystectomy. It has been shown that the pain reported by diclofenac group was statistically lower than that in placebo group in all intervals except at the time of recovery.

## 5. Discussion

It has been shown that NSAIDs reduce postoperative pain by blocking cyclo-oxygenase (COX) enzyme (2) and can be used as anesthetic drugs, solely or in combination with other opioids (4, 5). This randomized controlled clinical trial evaluated the efficacy of rectal diclofenac suppository in post laparoscopic cholecystectomy pain management. Preoperative characteristics of patients including age, sex, right upper quadrant and epigastric pain, and anorexia were not statistically different between treatment and placebo group. Moreover, intraoperative clinical findings including gall bladder adhesion and bleeding were not significantly different between them. Therefore, the results could be directly related to the effect of using diclofenac rectal suppository. In our study, the administration of postoperative opioid (pethedin) was limited to the patients whose VAS score was  $\geq 7$ . The results showed that mean pethedin consumption dose and the incidence of its administration were significantly lower in diclofenac group than that in placebo group. Opioids have critical side effects including nausea, vomiting, drowsiness, and respiratory depression especially in higher doses (6). Lack of need for opioids in diclofenac group would be able to decrease these side effects. This study demonstrated that the severity of post laparoscopic cholecystectomy pain was lower in the patients received 100 mg diclofenac rectal suppository than that in patients received placebo. This happened while the pain severity in placebo group might be masked by receiving higher dose of pethedin. Moreover, one of the side effects of using NSAIDs is increasing the risk of bleeding by inhibiting the effect of COX-1 in platelet aggregation and homeostasis (7). However, Intraoperative and postoperative bleeding was not statistically different between two groups. Most of the previous studies evaluated the efficacy of intramuscular diclofenac with regard to decreasing the post laparoscopic cholecystectomy pain. Fredman et al. (8) compared diclofenac, Ketrolac, and placebo groups. They showed that pain severity and need for opioid is lower in diclofenac and ketrolac group rather than in placebo one. In another study, Wilson et al. demonstrated the same results in addition to this fact that diclofenac would not increase the postoperative complications (9). Muscle damage and increasing the risk of bleeding and kidney injury are the important reported side effects of intramuscular administration of diclofenac (3). Therefore, a great need exists for finding other roots of administration. Limited number of studies evaluated the effect of other types of diclofenac on postoperative pain. Alessandri et al. demonstrated that subcutaneous diclofenac is effective in reducing postoperative pain (10). Moreover, Ng et al. compared the effect of suppository diclofenac to Intravenous parecoxib on postoperative pain. They showed that there is not statistically significant difference between them (11). Our study confirmed previous results and showed that 100 mg diclofenac rectal suppository at the time of recovery and three hours

later could reduce post laparoscopic cholecystectomy pain and also the need for opioid consumption. There-

fore, diclofenac rectal suppository would be a simple and effective alternative for this purpose.

**Table 1.** Baseline Characteristics of Patients

| Baseline Characteristics                | Diclofenac Group  | Placebo Group, No. (%) | P value |
|---|-------------------|------------------------|---------|
| <b>Sex, No. (%)</b>                     |                   |                        | 0.272   |
| Male                                    | 8 (36.4)          | 11 (50)                |         |
| Female                                  | 14 (63.6)         | 11 (50)                |         |
| <b>Age, Mean <math>\pm</math> SD</b>    | 47.05 $\pm$ 16.11 | 44.47 $\pm$ 17.72      | 0.658   |
| <b>Preoperative findings, No. (%)</b>   |                   |                        |         |
| Epigastric pain                         | 2 (9.1)           | 6 (27.3)               | 0.120   |
| RUQ <sup>a</sup> pain                   | 17 (77.3)         | 13 (59.1)              | 0.166   |
| Anorexia                                | 4 (18.2)          | 9 (40.9)               | 0.093   |
| <b>Intraoperative findings, No. (%)</b> |                   |                        |         |
| Gall bladder adhesion                   | 17 (77.3)         | 12 (54.5)              | 0.101   |
| Bleeding                                | 3 (13.6)          | 6 (27.3)               | 0.228   |

<sup>a</sup> Right Upper Quadrant

**Table 2.** Mean Visual Analogue Scale

|  | Diclofenac Group, Mean $\pm$ SD | Placebo Group, Mean $\pm$ SD | P value |
|--|---------------------------------|------------------------------|---------|
| <b>VAS<sup>a</sup> (Recovery time)</b> | 7.95 $\pm$ 0.899                | 8.32 $\pm$ 0.477             | 0.101   |
| <b>VAS (After 3 hours)</b>             | 6.59 $\pm$ 1.046                | 7.64 $\pm$ 0.685             | 0.013   |
| <b>VAS (After 6 hours)</b>             | 5.82 $\pm$ 1.468                | 7 $\pm$ 0.69                 | 0.001   |
| <b>VAS (After 12 hours)</b>            | 4.05 $\pm$ 1.588                | 5.91 $\pm$ 0.921             | < 0.001 |
| <b>VAS (After 24 hours)</b>            | 2.41 $\pm$ 1.501                | 4.59 $\pm$ 1.141             | < 0.001 |

<sup>a</sup> Visual Analogue Scale

## Acknowledgements

There is nothing to acknowledge.

## Authors' Contribution

The first author and corresponding author contributed 80% and the others 50%.

## Financial Disclosure

There is no financial disclosure.

## Funding Support

This research has been supported by Iran University of Medical Sciences.

## References

1. Fassoulaki A, Patris K, Sarantopoulos C, Hogan Q. The analgesic effect of gabapentin and mexiletine after breast surgery for cancer. *Anesth Analg*. 2002;**95**(4):985-91.
2. Dahl JB, Kehlet H. Non-steroidal anti-inflammatory drugs: rationale for use in severe postoperative pain. *Br J Anaesth*. 1991;**66**(6):703-12.
3. Power I. Muscle damage with diclofenac injections. *Anaesthesia*. 1992;**47**(5):451.
4. Hyllested M, Jones S, Pedersen JL, Kehlet H. Comparative effect of paracetamol, NSAIDs or their combination in postoperative pain management: a qualitative review. *Br J Anaesth*. 2002;**88**(2):199-214.
5. Rosenblum M, Weller RS, Conard PL, Falvey EA, Gross JB. Ibuprofen provides longer lasting analgesia than fentanyl after laparoscopic surgery. *Anesth Analg*. 1991;**73**(3):255-9.
6. Stein C. The control of pain in peripheral tissue by opioids. *N Engl J Med*. 1995;**332**(25):1685-90.
7. Sinatra R. Role of COX-2 inhibitors in the evolution of acute pain management. *J Pain Symptom Manage*. 2002;**24**(1 Suppl):S18-27.
8. Fredman B, Olsfanger D, Jedeikin R. A comparative study of ketorolac and diclofenac on post-laparoscopic cholecystectomy pain. *Eur J Anaesthesiol*. 1995;**12**(5):501-4.
9. Wilson YG, Rhodes M, Ahmed R, Daugherty M, Cawthorn SJ, Armstrong CP. Intramuscular diclofenac sodium for postoperative analgesia after laparoscopic cholecystectomy: a randomised, controlled trial. *Surg Laparosc Endosc*. 1994;**4**(5):340-4.
10. Alessandri F, Lijoi D, Mistrangelo E, Nicoletti A, Crosa M, Ragni N. Topical diclofenac patch for postoperative wound pain in laparoscopic gynecologic surgery: a randomized study. *J Minim Invasive Gynecol*. 2006;**13**(3):195-200.
11. Ng A, Swami A, Smith G, Emembolu J. Early analgesic effects of intravenous parecoxib and rectal diclofenac following laparoscopic sterilization: a double-blind, double-dummy randomized controlled trial. *J Opioid Manag*. 2008;**4**(1):49-53.